

Package: anthroplus (via r-universe)

November 12, 2024

Title Computation of the WHO 2007 References for School-Age Children and Adolescents (5 to 19 Years)

Version 1.0.0.9000

Description Provides WHO 2007 References for School-age Children and Adolescents (5 to 19 years) (z-scores) with confidence intervals and standard errors around the prevalence estimates, taking into account complex sample designs. More information on the methods is available online:
<<https://www.who.int/tools/growth-reference-data-for-5to19-years>>.

License GPL (>= 3)

Encoding UTF-8

Roxygen list(markdown = TRUE)

RoxygenNote 7.3.2

Depends R (>= 3.5.0)

Imports anthro (>= 1.0.0)

Suggests testthat (>= 3.0.0), survey

Config/testthat/edition 3

LazyData true

Collate 'data.R' 'zscores.R' 'prevalence.R'

URL <https://github.com/WorldHealthOrganization/anthroplus>

BugReports <https://github.com/WorldHealthOrganization/anthroplus/issues>

Config/pak/sysreqs make

Repository <https://worldhealthorganization.r-universe.dev>

RemoteUrl <https://github.com/worldhealthorganization/anthroplus>

RemoteRef HEAD

RemoteSha 799a7deaf7dff57f2abef5781f74b243f0aa6194

Contents

anthroplus_prevalence	2
anthroplus_zscores	4
Survey_WHO2007	6

Index	7
--------------	----------

anthroplus_prevalence *Compute prevalence estimates*

Description

Prevalence estimates according to the WHO recommended standard analysis: includes prevalence estimates with corresponding standard errors and confidence intervals, and z-score summary statistics (mean and standard deviation) with most common cut-offs describing the full index distribution (-3, -2, -1, +1, +2, +3), and at disaggregated levels for all available factors (age and sex).

Usage

```
anthroplus_prevalence(
  sex,
  age_in_months = NA_real_,
  oedema = "n",
  height_in_cm = NA_real_,
  weight_in_kg = NA_real_,
  sw = NULL,
  cluster = NULL,
  strata = NULL
)
```

Arguments

sex	A numeric or text variable containing gender information. If it is numeric, its values must be: 1 for males and 2 for females. If it is character, it must be "m" or "M" for males and "f" or "F" for females. No z-scores will be calculated if sex is missing.
age_in_months	A numeric variable containing age information; Age-related z-scores will NOT be calculated if age is missing.
oedema	The values of this character variable must be "n", "N" or "2" for non-oedema, and "y", "Y", "1" for oedema. Although it is highly recommended that this variable is provided by the survey, it is possible to run the analysis without specifying this variable. If unspecified, the default vector of all "n" with values considered as non-oedema is used. Missing values will be treated as non-oedema. For oedema, weight related z-scores are NOT calculated (set to missing), BUT they are treated as being < -3 SD in the weight-related indicator prevalence (anthroplus_prevalence) estimation.

height_in_cm	A numeric variable containing standing height information, which must be in centimeters. Height-related z-scores will not be calculated if missing.
weight_in_kg	A numeric variable containing body weight information, which must be in kilograms. Weight-related z-scores are not calculated if missing.
sw	An optional numeric vector containing the sampling weights. If NULL, no sampling weights are used.
cluster	An optional integer vector representing clusters. If the value is NULL this is treated as a survey without clusters. This is also the case if all values are equal, then it is assumed there are also no clusters.
strata	An optional integer vector representing strata. Pass NULL to indicate that there are no strata.

Details

In this function, all available (non-missing and non-flagged) z-score values are used for each indicator-specific prevalence estimation (standard analysis).

Note: the function temporarily sets the survey option `survey.lonely.psu` to "adjust" and then restores the original value. It is a wrapper around the survey package to compute estimates for the different groups (e.g. by age or sex).

If not all parameter values have equal length, parameter values will be repeated to match the maximum length.

Only cases with `age_in_months` between 60 (including) and 228 months (including) are used for the analysis. The rest will be ignored.

Value

Returns a data.frame with prevalence estimates for the various groups.

The output data frame includes prevalence estimates with corresponding standard errors and confidence intervals, and z-score summary statistics (mean and standard deviation) with most common cut-offs describing the full index distribution (-3, -2, -1, +1, +2, +3), and at disaggregated levels for all available factors.

The resulting columns are coded with a *prefix*, a *prevalence indicator* and a *suffix*:

Prefix:

HA Height-for-age

WA Weight-for-age

BMI Body-mass-index-for-age

Prevalence indicator:

_3 Prevalence corresponding to < -3 SD

_2 Prevalence corresponding to < -2 SD

_1 Prevalence corresponding to < -1 SD

1 Prevalence corresponding to > +1 SD

2 Prevalence corresponding to > +2 SD

3 Prevalence corresponding to $> +3$ SD**Suffix:**

_pop Weighted sample size
_unwpop Unweighted sample size
_r Mean/prevalence
_ll lower 95% confidence interval limit
_ul upper 95% confidence interval limit
_stdev Standard Deviation
_se Standard error

For example:

HA_r Height-for-age z-score mean
WA_stdev Weight-for-age z-score Standard Deviation
BMI_2_se Prevalence of BMI-for-age <-2 SD standard error
BMI_3_ll Prevalence of BMI-for-age <-3 SD lower 95% confidence interval limit

Note that weight-for-age results are NA for the groups "All" and the two "Sex" groups, as the indicator is only defined for age in months between 60 and 120.

Examples

```
set.seed(1)
prev <- anthroplus_prevalence(
  sex = c(1, 2),
  age_in_months = rpois(100, 100),
  height_in_cm = rnorm(100, 100, 10),
  weight_in_kg = rnorm(100, 40, 10)
)
prev[, c(1, 4, 5, 6)]
```

anthroplus_zscores *Compute z-scores for age 5 to 19*

Description

Compute z-scores for age 5 to 19

Usage

```
anthroplus_zscores(
  sex,
  age_in_months = NA_real_,
  oedema = NA_character_,
  height_in_cm = NA_real_,
  weight_in_kg = NA_real_
)
```

Arguments

sex	A numeric or text variable containing gender information. If it is numeric, its values must be: 1 for males and 2 for females. If it is character, it must be "m" or "M" for males and "f" or "F" for females. No z-scores will be calculated if sex is missing.
age_in_months	A numeric variable containing age information; Age-related z-scores will NOT be calculated if age is missing.
oedema	The values of this character variable must be "n", "N" or "2" for non-oedema, and "y", "Y", "1" for oedema. Although it is highly recommended that this variable is provided by the survey, it is possible to run the analysis without specifying this variable. If unspecified, the default vector of all "n" with values considered as non-oedema is used. Missing values will be treated as non-oedema. For oedema, weight related z-scores are NOT calculated (set to missing), BUT they are treated as being < -3 SD in the weight-related indicator prevalence (anthroplus_prevalence) estimation.
height_in_cm	A numeric variable containing standing height information, which must be in centimeters. Height-related z-scores will not be calculated if missing.
weight_in_kg	A numeric variable containing body weight information, which must be in kilograms. Weight-related z-scores are not calculated if missing.

Details

The following age cutoffs are used:

- Height-for-age age between 60 and 228 months inclusive
- Weight-for-age age between 60 and 120 months inclusive
- BMI-for-age age between 60 and 228 months inclusive

Value

A data.frame with three types of columns. Columns starting with a "c" are cleaned versions of the input arguments. Columns beginning with a "z" are the respective z-scores and columns prefixed by a "f" indicate if these z-scores are flagged (integers). The number of rows is given by the length of the input arguments.

The following columns are returned:

- age_in_months the input age in months
- csex standardized sex information
- coedema standardized oedema value
- cbmi BMI value based on weight/height
- zhfa Height-for-age z-score
- fhfa 1, if $\text{abs}(\text{zhfa}) > 6$
- zwfa Weight-for-age z-score
- fwfa 1, if $\text{zwfa} > 5$ or $\text{zwfa} < -6$
- zbfa BMI-for-age z-score
- fbfa 1, if $\text{abs}(\text{zbfa}) > 5$

Examples

```
anthroplus_zscores(  
  sex = c("f", "m"),  
  age_in_months = c(100, 110),  
  height_in_cm = c(100, 90),  
  weight_in_kg = c(30, 40)  
)
```

Survey_WHO2007

Sample Survey Data for the WHO 2007 References

Description

The dataset contains information of 933 children aged 61-228 months and it was created using several surveys; therefore, it does not represent any particular population.

Usage

Survey_WHO2007

Format

A data frame with 933 rows and 12 variables:

strata stratified sampling (integer)

cluster primary sampling unit (integer)

id child identification number (integer)

sex sex of the child (integer; 1=male/2=female)

dob date of birth (date)

dov date of visit (date)

agemons age in months (numeric)

weight child weight in kilograms (numeric)

height child height in centimeters (numeric)

sw child sample weight (numeric)

oedema presence of oedema (character; y=yes/n=no)

region geographical region (character; north/east/west/south)

Index

* datasets

Survey_WHO2007, [6](#)

anthroplus_prevalence, [2](#), [2](#), [5](#)

anthroplus_zscores, [4](#)

Survey_WHO2007, [6](#)